



## University of Arizona / Banner Health ECT Anesthesia Checklist

Confirm identity, consent, indication	
Review comorbidities & medications	<b>Follow BUMCP Time Out / Preprocedure Checklists</b>
NPO status verified	
Airway assessment complete	
IV access and baseline vitals	

<b>Setup (Routine)</b>	
Standard monitors (ECG, NIBP, SpO2, EtCO2)	
Suction, oxygen, BVM ready	
Airway equipment available	
ECT machine checked	

<b>Induction &amp; Paralysis (Critical)</b>	
Preoxygenate 100% O2	
Administer induction agent	
Administer succinylcholine	
Confirm mask ventilation	

Drug	Dose
Methohexital	0.75 – 1.0 mg/kg
Propofol	1.0 – 2.0 mg/kg
Etomidate	0.2 - 0.3 mg/kg
Ketamine	0.5 – 2.0 mg/kg
Remifentanyl	0.5 – 1.0 mcg/kg
Succinylcholine	1.0 mg/kg
Rocuronium	0.6 mg/kg
Sugammadex	16 mg/kg



<b>Seizure Phase (Critical)</b>	
Insert bite block	
Deliver stimulus	
Monitor seizure duration (≥20–25 sec)	
Support ventilation	

<b>Hemodynamics (Critical)</b>	
Treat bradycardia (atropine)	
Treat HTN/tachycardia (esmolol/labetalol)	

<b>Recovery (Routine)</b>	
Resume spontaneous ventilation	Lorem ipsum dolor sit amet, consectetur adipiscing elit,
Monitor for complications	Enter and sign PACU orders
Assess orientation & stability	Finalize anesthesia record and prep (if not already signed)